



eCOMMONS

Loyola University Chicago
Loyola eCommons

Bioinformatics Faculty Publications

Faculty Publications and Other Works by
Department

6-4-2020

Draft Genome Sequence of Streptococcus anginosus UMB0839, Isolated from the Female Urinary Tract

Thomas Scott
Loyola University Chicago

Taylor Miller-Ensminger
Loyola University Chicago

Adelina Voukadinova
Loyola University Chicago

Alan J. Wolfe
Loyola University Chicago, awolfe@luc.edu

Catherine Putoni
Loyola University Chicago, cputonti@luc.edu

Follow this and additional works at: https://ecommons.luc.edu/bioinformatics_facpub



Part of the [Bioinformatics Commons](#), and the [Biology Commons](#)

Recommended Citation

Scott, Thomas; Miller-Ensminger, Taylor; Voukadinova, Adelina; Wolfe, Alan J.; and Putoni, Catherine. Draft Genome Sequence of Streptococcus anginosus UMB0839, Isolated from the Female Urinary Tract. Microbiology Resource Announcements, 9, : , 2020. Retrieved from Loyola eCommons, Bioinformatics Faculty Publications, <http://dx.doi.org/10.1128/MRA.00411-20>

This Article is brought to you for free and open access by the Faculty Publications and Other Works by Department at Loyola eCommons. It has been accepted for inclusion in Bioinformatics Faculty Publications by an authorized administrator of Loyola eCommons. For more information, please contact ecommons@luc.edu.



This work is licensed under a [Creative Commons Attribution 4.0 License](#).
© Scott et al., 2020.



Draft Genome Sequence of *Streptococcus anginosus* UMB0839, Isolated from the Female Urinary Tract

Thomas Scott,^{a,b} Taylor Miller-Ensminger,^a Adelina Voukadinova,^a Alan J. Wolfe,^c  Catherine Putonti^{a,b,c,d}

^aBioinformatics Program, Loyola University Chicago, Chicago, Illinois, USA

^bDepartment of Biology, Loyola University Chicago, Chicago, Illinois, USA

^cDepartment of Microbiology and Immunology, Stritch School of Medicine, Loyola University Chicago, Maywood, Illinois, USA

^dDepartment of Computer Science, Loyola University Chicago, Chicago, Illinois, USA

ABSTRACT Here, we present the draft genome sequence of *Streptococcus anginosus* UMB0839, isolated from the female urinary tract. The total size of the genome is 2,104,786 bp assembled into 42 contigs with a GC content of 38.8% and 284× genome coverage.

Streptococcus anginosus is a member of the microbiota of the oral cavity, throat, and urinary tract. Previous studies on *S. anginosus* and other members of the *S. milleri* group (which also includes *S. intermedius* and *S. constellatus*) reported an association with purulent infection in both adults and children (1, 2). Compared to other members of the *S. milleri* group, *S. anginosus* is most abundant in the urogenital tract and is associated with urinary tract infections (1–3) and urgency urinary incontinence (4). Here, we sequenced an *S. anginosus* strain (UMB0839) isolated from a catheterized urine sample obtained from a pregnant woman.

The *Streptococcus anginosus* strain was collected as part of a prior institutional review board (IRB)-approved study (5), following the expanded quantitative urinary culture (EQUC) protocol, as previously described (6). The genus and species of the isolate were determined through matrix-assisted laser desorption ionization–time of flight (MALDI-TOF) mass spectrometry (6), after which it was stored at –80°C. The *S. anginosus* isolate was taken from the freezer and streaked onto a Columbia nalidixic acid (CNA) agar plate and incubated at 35°C with 5% CO₂ for 24 h. After incubation, a single colony was selected and inoculated in a liquid culture LB broth at 37°C with shaking for 24 h. The DNA was extracted using the Qiagen DNeasy blood and tissue kit, following the protocol for Gram-positive bacteria with slight modifications; 230 μl of lysis buffer (180 μl 20 mM Tris-Cl, 2 mM sodium EDTA, and 1.2% Triton X-100 and 50 μl of lysozyme) was used in step 2, and the incubation time in step 5 was reduced to 10 min. The DNA was quantified using the Qubit fluorometer and sent to the Microbial Genome Sequencing Center (MiGS) at the University of Pittsburgh for library preparation, using the Illumina Nextera kit, and sequencing, using the Illumina NextSeq 550 platform. Sequencing produced 1,565,124 pairs of 150-bp reads. The reads were trimmed using Sickel v1.33 (<https://github.com/najoshi/sickle>) and assembled using SPAdes v3.13.0 (7) with the “only-assembler” option for k values of 55, 77, 99, and 127. The contigs were annotated using the NCBI Prokaryotic Genome Annotation Pipeline (PGAP) v4.11 (8). The genome coverage was calculated using BBMap v38.47 (<https://sourceforge.net/projects/bbmap/>). Default parameters were used for all software unless previously stated.

The *S. anginosus* UMB0839 draft genome has a length of 2,104,786 bp and was assembled in 42 contigs. Its GC content is 38.8% with a coverage of 284×. The *N*₅₀ score for the genome is 117,003 bp. Annotation with PGAP identified 1,956 protein-coding

Citation Scott T, Miller-Ensminger T, Voukadinova A, Wolfe AJ, Putonti C. 2020. Draft genome sequence of *Streptococcus anginosus* UMB0839, isolated from the female urinary tract. Microbiol Resour Announc 9:e00411-20. <https://doi.org/10.1128/MRA.00411-20>.

Editor Kenneth M. Stedman, Portland State University

Copyright © 2020 Scott et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Catherine Putonti, cputonti@luc.edu.

Received 13 April 2020

Accepted 18 May 2020

Published 4 June 2020

genes alongside 47 tRNAs and 4 rRNA sequences (2 5S, 1 16S, and 1 23S). Future work will help further our understanding of the role that *S. anginosus* plays in the urogenital microbiota, especially its tendency to cause infection. Moreover, the addition of this strain to the several *S. anginosus* strains already sequenced will help to clarify the genetic characteristics of *S. milleri* bacteria and their identification.

Data availability. This whole-genome shotgun project has been deposited in GenBank under the accession no. [JAAUWM000000000](https://www.ncbi.nlm.nih.gov/nuclseq/JAAUWM000000000). The version described in this paper is the first version, JAAUWM010000000. The raw sequencing reads have been deposited in the SRA under the accession no. [SRR11441025](https://www.ncbi.nlm.nih.gov/sra/SRR11441025).

ACKNOWLEDGMENTS

This work was conducted as part of the Bacterial Genomics course at Loyola University Chicago's Department of Biology. For prior patient recruitment, we acknowledge the Loyola Urinary Education and Research Collaborative (LUEREC) and the patients who provided the samples for this study.

REFERENCES

1. Whiley RA, Beighton D, Winstanley TG, Fraser HY, Hardie JM. 1992. *Streptococcus intermedius*, *Streptococcus constellatus*, and *Streptococcus anginosus* (the *Streptococcus milleri* group): association with different body sites and clinical infections. *J Clin Microbiol* 30:243–244. <https://doi.org/10.1128/JCM.30.1.243-244.1992>.
2. Furuichi M, Horikoshi Y. 2018. Sites of infection associated with *Streptococcus anginosus* group among children. *J Infect Chemother* 24:99–102. <https://doi.org/10.1016/j.jiac.2017.09.011>.
3. Jacobs JA, Pietersen HG, Stobberingh EE, Soeters PB. 1995. *Streptococcus anginosus*, *Streptococcus constellatus* and *Streptococcus intermedius* clinical relevance, hemolytic and serologic characteristics. *Am J Clin Pathol* 104:547–553. <https://doi.org/10.1093/ajcp/104.5.547>.
4. Pearce MM, Hilt EE, Rosenfeld AB, Zilliox MJ, Thomas-White K, Fok C, Kliethermes S, Schreckenberger PC, Brubaker L, Gai X, Wolfe AJ. 2014. The female urinary microbiome: a comparison of women with and without urgency urinary incontinence. *mBio* 5:e01283-14. <https://doi.org/10.1128/mBio.01283-14>.
5. Jacobs KM, Thomas-White KJ, Hilt EE, Wolfe AJ, Waters TP. 2017. Microorganisms identified in the maternal bladder: discovery of the maternal bladder microbiota. *AJP Rep* 7:e188–e196. <https://doi.org/10.1055/s-0037-1606860>.
6. Hilt EE, McKinley K, Pearce MM, Rosenfeld AB, Zilliox MJ, Mueller ER, Brubaker L, Gai X, Wolfe AJ, Schreckenberger PC. 2014. Urine is not sterile: use of enhanced urine culture techniques to detect resident bacterial flora in the adult female bladder. *J Clin Microbiol* 52:871–876. <https://doi.org/10.1128/JCM.02876-13>.
7. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
8. Tatusova T, DiCuccio M, Badretdin A, Chetverin V, Nawrocki EP, Zaslavsky L, Lomsadze A, Pruitt KD, Borodovsky M, Ostell J. 2016. NCBI Prokaryotic Genome Annotation Pipeline. *Nucleic Acids Res* 44:6614–6624. <https://doi.org/10.1093/nar/gkw569>.